

P.O. Bur: 48577, DUBAI, U.A.E. Tel: 04:267 8886, Fan: 04:267 8856 P.B. Hart 457. SHARJAH, U.A.E.

E-mail: Info@pulokhahospitols.com • Webelle: www.rulekhahospitals.com

ميرب ١٨٨٧٧ نيني ( ع.ب -1 476 AAA . COL -5 479 AV71 . - EEA عرب ١٠٧٠ فلمارقة ( ١ وم 

# CORONARY **ANGIOPLASTY**



1. What is Coronary angioplasty?

Coronary angioplasty (PCI or PTCA) is a technique for treating coronary artery blockage.

## 2. How is coronary angioplasty done?

A fine flexible tube with a small inflatable balloon at its top is pierced in to the artery in either the groin or arm. This balloon is positioned across the block in your heart vessel under X ray screening. The balloon will be gently inflated so that it substitutes the fatty tissue of the removed artery.

### 3. What is a stent?

After Angioplasty usually stenting is done to prevent the recoil and re block of the heart blood vessel. Here a catheter containing stent- a small tube of stainless steel mesh is inserted across the block and inflated.

#### 4. What are the types of stents?

There are two types of stents:

- Base metal stent made up of stainless steel or cobalt chromium.
- Drug eluding stent- here the stent is coated with a drug which is released slowly over 6 months.

#### 5. What is the advantage of drug eluding stent?

Drug eluding stent prevents the chance of re blockage. The chance of re blockage is because of 5% after drug eluding stent is implanted.

#### 6. What happens after the procedure?

After the procedure the catheter is removed. There might be small amount of bleeding when the catheter is taken out. After wards s compression bandage or a plus angroseal is put to stop any bleeding. You will be kept in the hospital for 2 days for monitoring.

## 7. What precautions should I take after the procedure?

- After the discharge watch for any bleeding or swelling in the catheter insertion site. If you have any bleeding or swelling immediately report to the Doctor or to the Hospital.
- If you have any signs of infection, seen as redness, pus or fever then you should visit the Doctor.
- In case you develop any chest pain after going home.
  - Stop what you are doing.
  - Sit down and rest.
  - > Take your GTN spray or tablets.

Take the GTN spray as your Doctor or nurse has instructed you. The pain should subside with in a few minutes. If it does not get relieved take another GTN and call emergency immediately.